

No. of studies	Study design	Sample size	Celebrate & Protect (% vaccine uptake)	Control (% vaccine uptake)	Risk of bias	Inconsistency	Indirectness	Imprecision	Quality
<b>12 months of age (Diphtheria, tetanus, pertussis, polio, Hib)</b>									
1 (Gibson 2014)	Non-randomised cluster	Not reported	85.7%	88.0%	Critical <sup>2</sup>	N/A <sup>3</sup>	Not serious	N/A <sup>4</sup>	Very low
<b>12 months of age (Meningitis C)</b>									
1 (Gibson 2014)	Non-randomised cluster	Not reported	89.5%	92.1%	Critical <sup>2</sup>	N/A <sup>3</sup>	Not serious	N/A <sup>4</sup>	Very low

No. of studies	Study design	Sample size	Celebrate & Protect (% vaccine uptake)	Control (% vaccine uptake)	Risk of bias	Inconsistency	Indirectness	Imprecision	Quality
<b>24 months of age (Hib and Meningitis C)</b>									
1 (Gibson 2014)	Non-randomised cluster	Not reported	84.4%	83.3%	Critical <sup>2</sup>	N/A <sup>3</sup>	Not serious	N/A <sup>4</sup>	Very low
<b>24 months of age (MMR 1<sup>st</sup> dose)</b>									
1 (Gibson 2014)	Non-randomised cluster	Not reported	86.3%	84.5%	Critical <sup>2</sup>	N/A <sup>3</sup>	Not serious	N/A <sup>4</sup>	Very low
<b>5 years of age (MMR 1<sup>st</sup> dose)</b>									
1 (Gibson 2014)	Non-randomised cluster	Not reported	91.3%	87.7%	Critical <sup>2</sup>	N/A <sup>3</sup>	Not serious	N/A <sup>4</sup>	Very low
<b>5 years of age (MMR 2<sup>nd</sup> dose)</b>									
1 (Gibson 2014)	Non-randomised cluster	Not reported	80.5%	78.7%	Critical <sup>2</sup>	N/A <sup>3</sup>	Not serious	N/A <sup>4</sup>	Very low
<ol style="list-style-type: none"> <li>1. Modified GRADE table. Study did not report number of participants so not possible to calculate risk ratios or absolute risk. Quality assessment based on risk of bias and indirectness.</li> <li>2. Single study at critical risk of bias. Quality of the outcome downgraded three times.</li> <li>3. Single study. Inconsistency not applicable.</li> <li>4. Not possible to calculate risk ratios so imprecision could not be assessed.</li> </ol>									